

First record of *Osmundea oederi* (Gunnerus) G. Furnari comb. nov. (Rhodomelaceae, Rhodophyta) from the Mediterranean Sea

Donatella SERIO^{a*}, Antonella PETROCELLI^b, Mario CORMACI^a,
Ester CECERE^b and Giovanni FURNARI^a

^aDepartment of Botany of the University of Catania, via A. Longo 19,
95125 Catania, Italy

^bInstitute for Marine Coastal Environment, Talassografico "A. Cerruti" –
C.N.R., Via Roma 3, 74100 Taranto, Italy

(Received 18 July 2007, accepted 11 January 2008)

Abstract – The first documented occurrence of *Osmundea oederi* (Gunnerus) G. Furnari comb. nov. in the Mediterranean Sea is reported from the Gulf of Taranto (Italy). A comparison with the morphologically similar species *O. truncata* and other species of *Osmundea* with compressed thalli and secondary pit-connections between epidermal cells occurring in the Mediterranean Sea, is presented. The conspecificity of the Caribbean *Laurencia bolivarii* with *O. oederi* is also proposed.

Mediterranean Sea / *Osmundea oederi* / Rhodomelaceae / Rhodophyta / Taranto

Résumé – Premier report d'*Osmundea oederi* (Gunnerus) G. Furnari comb. nov. en mer Méditerranée. La présence d'*Osmundea oederi* (Gunnerus) G. Furnari comb. nov. est signalée pour la première fois en Méditerranée dans le Golfe de Tarante (Italie). *O. oederi* est comparé avec *O. truncata*, espèce morphologiquement voisine, et avec d'autres espèces d'*Osmundea* présentes en Méditerranée et pourvues de thalles comprimés et de synapses secondaires entre les cellules de l'épiderme. *Laurencia bolivarii* des Caraïbes est conspécifique d'*O. oederi*.

Mer Méditerranée / *Osmundea oederi* / Rhodomelaceae / Rhodophyta / Golf de Tarante

INTRODUCTION

In the Mediterranean Sea, the *Laurencia* complex is represented by 20 taxa at specific and infraspecific levels (Table 1), belonging to the genera *Laurencia* (Hudson) J.V. Lamouroux, *Osmundea* Stackhouse, *Chondrophycus* (Tokida & Saito) Garbary & J. Harper and *Palisada* (Yamada) K.W. Nam.

* Correspondence and reprints: d.serio@unict.it
Communicating editor: Frederik Leliaert

Table 1. Taxa of the *Laurencia* complex recorded to date from the Mediterranean Sea.

Taxon	References
<i>Chondrophycus patentirameus</i> (Montagne) K.W. Nam	Furnari <i>et al.</i> (2001)
<i>Chondrophycus tenerrimus</i> (Cremades) G. Furnari, Boisset	Furnari <i>et al.</i> (2002)
<i>Chondrophycus thuyoides</i> (Kützing) G. Furnari	Furnari <i>et al.</i> (2001)
<i>Laurencia caduciramulosa</i> Masuda & Kawaguchi	Furnari <i>et al.</i> (2001)
<i>Laurencia chondrioides</i> Børgesen	Furnari <i>et al.</i> (2001)
<i>Laurencia epiphylla</i> Boisset & Lino	Furnari <i>et al.</i> (2001)
<i>Laurencia glandulifera</i> (Kützing) Kützing	Furnari <i>et al.</i> (2001)
<i>Laurencia intricata</i> J.V. Lamouroux	Furnari <i>et al.</i> (2001)
<i>Laurencia majuscula</i> (Harvey) Lucas	Furnari <i>et al.</i> (2001)
<i>Laurencia microcladia</i> Kützing	Furnari <i>et al.</i> (2001)
<i>Laurencia minuta</i> Vandermeulen, Garbary & Guiry subsp. <i>scammacae</i> G. Furnari & Cormaci	Furnari <i>et al.</i> (2001)
<i>Laurencia obtusa</i> (Hudson) J.V. Lamouroux	Furnari <i>et al.</i> (2001)
<i>Laurencia pyramidalis</i> Bory ex Kützing	Serio <i>et al.</i> (2004)
<i>Osmundea maggsiana</i> Serio, Cormaci & G. Furnari	Furnari <i>et al.</i> (2001)
<i>Osmundea pelagiensis</i> G. Furnari	Furnari <i>et al.</i> (2001)
<i>Osmundea pelagosae</i> (Schiffner) K.W. Nam	Furnari <i>et al.</i> (2001)
<i>Osmundea pinnatifida</i> (Hudson) Stackhouse	Furnari <i>et al.</i> (2001)
<i>Osmundea truncata</i> (Kützing) K.W. Nam & Maggs	Furnari <i>et al.</i> (2001)
<i>Osmundea verlaquei</i> G. Furnari	Furnari <i>et al.</i> (2001)
<i>Palisada papillosa</i> (C. Agardh) K.W. Nam	Furnari <i>et al.</i> [2001 as <i>Chondrophycus papillosus</i> (C. Agardh) Garbary & J. Harper]

Phylogenetic relationships within the *Laurencia* complex were studied by Nam (2006). The only Mediterranean species in the 2006 study, *Chondrophycus papillosus*, subsequently was ascribed to the newly validated genus *Palisada* (Nam, 2007).

In the course of a floristic study of the Mar Piccolo of Taranto (southern Italy, Ionian Sea), tetrasporic, male and female specimens of *Osmundea oederi* (Gunnerus) G. Furnari comb. nov. were found. This species was considered by Athanasiadis (1996, as *O. ramosissima* Athanasiadis *nom. illeg.*; see below) to be conspecific with *O. truncata*, a species widely distributed throughout the Mediterranean Sea. Nam *et al.* (2000) demonstrated, however, that *O. truncata* is distinct from *O. ramosissima* (= *O. oederi*). According to Nam *et al.* (2000), *O. ramosissima* (= *O. oederi*) occurred only in the NE Atlantic (France, Britain, Ireland, Helgoland and West Norway), and hence all the Mediterranean records of *O. ramosissima* (= *O. oederi*) were considered to represent *O. truncata* (Furnari *et al.*, 2001; Furnari *et al.*, 2003). Our present finding represents the first documented record of *Osmundea oederi* in the Mediterranean Sea.

MATERIAL AND METHODS

Investigations were carried out on tetrasporic, male and cystocarpic specimens collected at 0.5 m depth in January and February 2007 in the Mar Piccolo of Taranto (Fig. 1) by Cecere and Petrocelli. Observations were made on specimens-preserved in a 4% formaldehyde-seawater solution. Herbarium specimens are held at the Department of Botany of the University of Catania (CAT 2064 and 2690) and at the Herbarium of the Istituto Sperimentale Talassografico of Taranto (TAR 219 and 221). The following additional herbarium specimens were also examined: male and tetrasporic thalli (CAT 1151 and 1153, respectively) collected on 14/x/1987 in the Mar Piccolo of Taranto by Cecere and labelled as *O. truncata*.

Herbarium abbreviations follow Holmgren *et al.* (1990). For microscopic observations, some specimens were stained with 1% aqueous aniline blue acidified with dilute HCl, which enhances pit connections. Photographs were made by a Nikon D1 digital camera and mounted using Adobe Photoshop 6.0.

RESULTS

Thalli are epilithic, brown to reddish brown, up to 10 cm high, soft in texture, compressed throughout, except near the base of axes and in ultimate




Fig. 1. Location of the collection site (arrow).


branchlets which are terete (Fig. 2). Axes, up to 1.5 mm in diameter, arise from a discoid holdfast, are irregularly ramified and bear distichous, alternately or suboppositely arranged branches. In surface view, epidermal cells, with secondary pit-connections (Fig. 3), are elongate in basal to median parts of the thallus (50-60 µm long × 10-13 µm wide) and more or less isodiametric (25-30 µm long × 15-25 µm wide) near the apex. In transverse section, they are isodiametric and not palisade-like (Fig. 4). Medullary cells do not show lenticular thickenings (Fig. 4). Two pericentral cells are produced from each axial cell. Tetransporangia are produced randomly (Fig. 5) from epidermal cells and are cut off from the mother cells laterally. The two presporangial cover cells are aligned parallel to the stichidial axis in surface view. Male reproductive structures are of the filament-type, with spermatangial branches unramified or 1-2 alternately ramified. They usually terminate in a cluster of large vesicular sterile cells, 60-80 (100) µm high × 25-35 µm wide (Fig. 6), and are inserted in cup-like depressions, slightly sunken, and located at the bifurcation of branchlets (Fig. 7). Cystocarps, distributed in the middle to subapical parts of branches, are ovoid, 600-650 µm high × 620-700 µm wide, with non protuberant ostioles (Fig. 8).

Nomenclature

Athanasiadis (1996: 119) proposed the new combination *Osmundea ramosissima* (Oeder) Athanasiadis (basionym *Fucus ramosissimus* Oeder, 1766: pl. 276), on the erroneous premise that Oeder's "*Fucus, ramosissimus, ramis vagis approximatis teretibus, vesicis solidis, sparsis, caulinis*" is a validly published name. Lyngbye (1819) and C. Agardh (1822) also made such an incorrect interpretation, transforming Oeder's polynomial to the binomial *Fucus ramosissimus*. Lyngbye (1819: 213) treated *Fucus ramosissimus* as a synonym of *Gelidium pinnatifidum* β *angustum* listing the disposition of taxa in Oeder's (1766) *Flora Danica*; and on p. 40 he only mentioned Oeder's pl. 276 but not a name. C. Agardh (1822: 239) treated *Fucus ramosissimus* as a synonym of *Chondria pinnatifida*. Therefore, according to Art. 34.1 of ICBN-Vienna Code (McNeill *et al.*, 2006), neither the Lyngbye nor the Agardh transformed binomials are validly published, because they are merely cited as synonyms.

By contrast, Gunnerus (1772), Hornemann (1827), Lange (1887) as well as Nam *et al.* (2000), McIvor *et al.*, (2002) and P.C. Silva (personal communication) have correctly interpreted Oeder's "*Fucus, ramosissimus, ramis vagis approximatis teretibus, vesicis solidis, sparsis, caulinis*" as a polynomial and consequently not validly published according to Art. 23.1 of ICBN-Vienna Code (McNeill *et al.*, 2006). On this basis, Nam *et al.* (2000) stated that *O. ramosissima* should be considered a new species described by Athanasiadis. Nam *et al.* (2000) were unaware, however, that *O. ramosissima* Athanasiadis is a superfluous name according to Arts 52.1 and 52.2 of ICBN-Vienna Code (McNeill *et al.*, 2006) because the names of earlier validly described species are cited as synonyms by Athanasiadis (1996).

Later, McIvor *et al.* (2002) stated that *Fucus oederi* Gunnerus (1772) is the earliest available name for "*O. ramosissima*". However, the combination in *Osmundea* could not be made in that paper, which appeared in the online journal *Costancea*. In fact, Art. 29 of ICBN (Greuter *et al.*, 2000; McNeill *et al.*, 2006) specifically states that online publication is not effective for nomenclatural purposes. Therefore, the following new combination is proposed here:



Figs 2-8. *Osmundea oederi*. 2. Habit. Scale bar = 2 cm. 3. Epidermal cells in surface view showing secondary pit connections. Scale bar = 50 µm. 4. Transverse section of an axis showing medullary cells without lenticular thickenings. Scale bar = 100 µm. 5. Detail of a stichidial branchlet showing randomly disposed tetrasporangia. Scale bar = 2 mm. 6. Spermatangial branch terminating in a cluster of elongate vesicular sterile cells. Scale bar = 50 µm. 7. Male gametophyte with cup-like spermatangial depressions (arrows). Scale bar = 2 mm. 8. Female gametophyte with cystocarps (arrows). Scale bar = 2 mm.

***Osmundea oederi* (Gunnerus) G. Furnari, comb. nov.**

Basionym: *Fucus oederi* Gunnerus, *Flora norvegica*...2: 100 (1772).

Lectotype: "Fucus ramosissimus" Oeder, *Flora Danica*: plate 276 (1776).

Epitype: "Fucus ramosissimus" collected by M. N. Blytt, 1833 at Stavanger (Norway), designated by Nam *et al.* (2000: 767 and fig. 30); epitype deposited in O (the Botanical Museum of Oslo).

Synonym: *Osmundea ramosissima* Athanasiadis, *Taxonomisk litt...*:119 (1996)
nom. illeg.

DISCUSSION

Most of the characters of the Italian thalli are in agreement with the description of *Osmundea oederi* as provided by Nam *et al.* (2000, as *O. ramosissima*) for NE Atlantic specimens as well as with the epitype, except for minor features such as the size of vesicular cells in male structures [60-80 (100) µm high × 25-35 µm wide in Mediterranean specimens vs. 40-50 µm × 30-40 µm in NE Atlantic specimens] and of cystocarps (600-650 µm high × 620-700 µm wide in Mediterranean thalli vs 800-1300 µm × 800-1300 µm in NE Atlantic specimens). This species is morphologically very similar to *O. truncata*, from which it is almost indistinguishable when sterile. *O. oederi* differs from *O. truncata* by: i. the absence of lenticular thickenings in the medullary cells (abundantly present in *O. truncata*); ii. the absence of refractive needle-like inclusions in medullary cells (present in *O. truncata*); iii. cystocarps ovoid without protuberant ostioles (urceolate with protuberant ostioles in *O. truncata*); iv. spermatangial branches unramified or 1-2 alternately ramified and usually terminating in a cluster of vesicular sterile cells (spermatangial branches always unramified and terminating in a single large vesicular sterile cell in *O. truncata*).

As mentioned above, to date *Osmundea oederi* was only recorded from Atlantic France (Roscoff), Britain, Ireland, Helgoland and western Norway (Stavanger) (Nam *et al.*, 2000: 761, as *O. ramosissima*), so our finding represents the first documented record of the species from the Mediterranean Sea. However, it should be noted that since the above mentioned re-examined herbarium specimens labelled as *O. truncata* actually belong to *O. oederi*, the species has been present in the Mar Piccolo of Taranto since 1987. Moreover, in agreement with Nam *et al.* (2000), it cannot be excluded that the species recorded from the Gulf of Naples by Falkenberg (1901) as *Laurencia pinnatifida* (Hudson) J.V. Lamouroux, could be referred to as *O. oederi* on the basis of Falkenberg's illustrations (pl. 23: figs 21, 22, 33).

To date, only two other species of *Osmundea* with secondary pit connections between cortical cells are known from the Mediterranean Sea: *O. pelagosae* and *O. verlaquei*. *O. pelagosae* differs from *O. oederi* in having: medullary cells with lenticular thickenings (absent in *O. oederi*), deep and urn-shaped spermatangial receptacles with spermatangial branches only unramified and terminating with a single vesicular sterile cell (cup-like and slightly sunken spermatangial receptacles with spermatangial branches unramified and/or ramified terminating with a cluster of several vesicular sterile cell in *O. oederi*) (Table 2). Moreover, *O. pelagosae* occurs throughout the sublittoral zone (Furnari *et al.*, 2001) while *O. oederi* grows in the midlittoral zone (rarely in the subtidal

Table 2. Comparison of Mediterranean species of *Osmundea* with compressed thallus and secondary pit-connections between epidermal cells.

Species	Vegetative structures			Male structures		Female structure	Habitat	Geographical distribution	References
	Basal system	Lenticular thickenings	Spermatangial branch	Ending of spermatangial branch	Shape of spermatangial depression				
<i>Osmundea pelagosa</i> e	discoid holdfast	present	unramified	one elongate cell	urn-shaped	ovoid without protuberant ostiole	epiphytic, throughout the sublittoral zone.	Mediterranean	Furnari & Serio, 1993a [as <i>L. pelagosae</i> (Schiffner) Ercgovic]
<i>Osmundea oederi</i>	discoid holdfast	absent	unramified or ramified	one or a cluster of spherical to elongate apical cells	cup-like slightly sunken	ovoid without protuberant ostiole	epiphytic or epiphytic in the midlitoral zone (rarely to 7 m depth)	NE Atlantic; Caribbean Sea (as <i>L. boliviari</i>); Mediterranean	Nam et al., 2000; this study
<i>Osmundea truncata</i>	discoid holdfast	present	unramified	one spherical cell	cup-like, shallow and broad	urceolate with protuberant ostiole	epiphytic or epiphytic in mid- to upper sublittoral zones	NE Atlantic; Mediterranean	Furnari & Serio, 1993b [as <i>L. truncata</i> Küizing]; Nam et al., 1994; Nam et al., 2000
<i>Osmundea verlaquei</i>	spreading crust	absent	unramified or ramified	one or a cluster of elongate apical cells	cup-like slightly sunken	urceolate with more or less protuberant ostiole	epiphytic in the lower midlitoral zone	Mediterranean	Cormaci et al., 1994 (as <i>L. verlaquei</i> Cormaci et al.); this study

zone to 7 m depth) (Nam *et al.*, 2000). *O. verlaquei* differs from *O. oederi* in having crustose spreading basal system (discoid holdfast in *O. oederi*) and in urceolate cystocarps with protuberant ostiole (ovoid without protuberant ostiole in *O. oederi*) (Table 2).

Recently, García (2006) examined specimens of *Laurencia bolivarii* N. Rodríguez de Ríos (1981: 4-7, figs 6-9), including the type material from the coast of Venezuela. She demonstrated that they belonged to the genus *Osmundea*, and proposed the new combination *Osmundea bolivarii* (Rodríguez) M.García-Ortiz. However, this new combination is invalid, since the author gave neither the basionym nor its full and direct reference as required by Art. 33.4 of ICBN (McNeill *et al.*, 2006). Moreover, from both the description and iconography, the species seems to correspond to *O. oederi*. In particular, the following characters are shared by the two species: habit; branching pattern; presence of secondary pit-connections between epidermal cell; absence of lenticular thickenings in the wall of medullary cells; spermatangial branches unramified and usually terminating in a cluster of sterile cells, inserted in cup-like depressions, slightly sunken, located at the bifurcation of branchlets; cystocarps ovoid with not protuberant ostiole. Therefore, we conclude that ***Laurencia bolivarii* is a heterotypic synonym of *Osmundea oederi*.**

Finally, the previous known distribution of *Osmundea oederi* suggested a restricted cold water affinity even though the southernmost and northernmost distribution of the species is uncertain due to taxonomic confusion. However, its occurrence in both Venezuela (Caribbean Sea) and the Mar Piccolo of Taranto, a basin with a relatively high sea water temperature, indicates that the species shows a wide range of tolerance to this environmental factor. Moreover, even though due to its occurrence in the Mar Piccolo of Taranto, a basin subject to aquaculture and commercial fishing, *O. oederi* could be considered as an alien species in the Mediterranean Sea, past confusion with *O. truncata* also supports the hypothesis that *O. oederi* could have a wider distribution in the Mediterranean Sea into which it came by natural way.

Acknowledgements. We are grateful to Dr C.A. Maggs, Dr P.C. Silva and W.J. Woelkerling for useful discussion on nomenclature. Cecere and Petrocelli acknowledge the support of the Research Project VECTOR ("VulnErabilità delle Coste e degli ecosistemi marini italiani ai cambiamenti climatici e loro ruolo nei cicli del carbonio mediterraneo") financed by an Integrative Special Fund for the Research (FISR) – call 2001. The other authors acknowledge the financial support of the University of Catania.

REFERENCES

- AGARDH C., 1822 — *Species algarum*. I, part 2. Lund: VIII+169-398.
- ALABISO G., CANNALIRE M., GHIONDA D., MILILLO M., LEONE G. & CACIORGNA O., 1997 — Particulate matter and chemical-physical conditions of an inner sea: the Mar Piccolo in Taranto. A new statistical approach. *Marine chemistry* 58: 373-338.
- ATHANASIADIS A., 1996 — *Taxonomisk litteratur och biogeografi av Skandinaviska rödalger och brunalger*. Göteborg, Algologia, 280 p.
- CORMACI M., FURNARI G. & SERIO D., 1994 — Two new species of *Laurencia* (Ceramiales, Rhodophyta) from the Mediterranean Sea: *Laurencia pelagiensis* sp. nov. and *L. verlaquei* sp. nov. *The Japanese journal of phycology* 42: 365-375.
- FALKENBERG P., 1901 — Die Rhodomelaceen des Golfs von Neapel und der angrenzenden Meeres-Abschnitte. *Fauna und Flora des Golfs von Neapel, Monographie* 26. Berlin. XVI+754 p., 10 figs, 24 pls.

- FURNARI G. & SERIO D., 1993a — The reproductive structures of the Mediterranean alga *Laurencia pelagosa* (Ceramiales, Rhodophyta). *European journal of phycology* 28: 141-143.
- FURNARI G. & SERIO D., 1993b — The distinction of *Laurencia truncata* (Ceramiales, Rhodophyta) in the Mediterranean Sea from *Laurencia pinnatifida*. *Phycologia* 32: 367-372.
- FURNARI G., CORMACI M. & SERIO D., 2001 — The *Laurencia* complex (Rhodophyta, Rhodomelaceae) in the Mediterranean Sea: an overview. *Cryptogamie, Algologie*, 22: 331-373.
- FURNARI G., BOISSET F., CORMACI M. & SERIO D., 2002 — The characterization of *Chondrophycus tenerimus* (Cremades) comb. nov. (Ceramiales, Rhodophyta), a species often misidentified as *C. papillosum* (C. Agardh) Garbary et J. Harper in the Mediterranean Sea. *Cryptogamie, Algologie* 23: 223-235.
- FURNARI G., GIACCONE G., CORMACI M., ALONGI G. & SERIO D., 2003 — Biodiversità marina delle coste italiane: catalogo del macrofitobenthos. *Biologia marina Mediterranea* 10: 3-483.
- GARCÍA M., 2006 — *Osmundea bolivarii* (Rodríguez) M. García-Ortiz comb. nov. (Rhodomelaceae, Rhodophyta). *Acta botanica Venezolana* 29: 17-26.
- GREUTER W., McNEILL J., BARRIE F.R., BURDET H.M., DEMOULIN V., FILGUEIRAS T.S., NICOLSON D.H., SILVA P.C., SKOG J.E., TURLAND N.J. (Members) & HAWKSWORTH D.L. (Secretary of the Editorial Committee), 2000 — *International Code of Botanical Nomenclature (Saint Louis Code)*. Adopted by the Sixteenth International Botanical Congress, Saint Louis, Missouri, July-August 1999. [Regnum vegetabile, 131], xviii+389 p.
- GUNNERUS J.E., 1772 — *Flora norvegica... pars posterior*. Hafniae [Copenhagen], 148 p.
- HOLMGREN P.K., HOLMGREN N.H. & BARNETT L.C., 1990 — *Index Herbariorum. Part I. The herbaria of the World*. New York Botanical Garden, Bronx, N.Y. [Regnum Vegetabile, 120], x+693 p.
- HORNEMANN J.W., 1827 — *Nomenclatura Floraen danicae...* Hafniae [Copenhagen], 214 p.
- LANGE J.M.C., 1887 — *Nomenclator "Floraen danicae..."* Hauniae [Copenhagen], 354 p.
- LYNGBYE H.C., 1819 — *Tentamen hydrophytologiae danicae...* Hafniae [Copenhagen], 248 p.
- MCIVOR L., MAGGS C.A., GUIRY M.D. & HOMMERSAND M.H., 2002 — Phylogenetic analysis of the geographically disjunct genus *Osmundea* Stackhouse (Rhodomelaceae, Rhodophyta). *Constancea* 83 (<http://ucjeps.berkeley.edu/constancea/>).
- McNEILL J., BARRIE F.R., BURDET H.M., DEMOULINE V., HAWKSWORTH D.L., MARHOLD K., NICOLSON D.H., PRADO J., SILVA P.C., SKOG J.E., WIERSEMA J.H. & TURLAND N.J., 2006 — *International Code of Botanical Nomenclature (Vienna Code)* adopted by the Seventeenth International Botanical Congress Vienna, Austria, July 2005. Liechtenstein, A.R.G. Gantner Verlag, [i-iv], v-xviii + 1-568 p.
- NAM K.W., MAGGS C.A. & GARBARY D.J., 1994 — Resurrection of the genus *Osmundea* with an emendation of the delineation of *Laurencia* (Ceramiales, Rhodophyta). *Phycologia* 33: 384-395.
- NAM K.W., MAGGS C.A., MCIVOR L. & STANHOPE M.J., 2000 — Taxonomy and phylogeny of *Osmundea* (Rhodomelaceae, Rhodophyta) in Atlantic Europe. *Journal of phycology* 36: 759-772.
- NAM K.W., 2006 — Phylogenetic re-evaluation of the *Laurencia* complex (Rhodophyta) with a description of *L. succulenta* sp. nov. from Korea. *Journal of applied phycology* 18: 679-697.
- NAM K.W., 2007 — Validation of the generic name *Palisada* (Rhodomelaceae, Rhodophyta). *Algae* 22: 53-55.
- OEDER G.C., 1766 — *Icones plantarum sponte nascentium in regnis Daniae et Norvegiae...[Flora danica]*, vol. 2, fasc. 5. Hafniae [Copenhagen], tabs 241-300.
- RODRÍGUEZ DE RIOS N., 1981 — Dos especies nüevas de *Laurencia* (Rhodophyta, Ceramiales). *Ernstia* 2: 1-11.
- SERIO D., FURNARI G. & CORMACI M., 2004 — On the occurrence of *Laurencia pyramidalis* Bory ex Kützing (Rhodophyta, Rhodomelaceae) in the Mediterranean Sea. *Cryptogamie, Algologie* 25: 329-336.